

Jurnal Ilmiah
FARMASI

(Scientific Journal of Pharmacy)



JURNAL ILMIAH FARMASI
(SCIENTIFIC JOURNAL OF PHARMACY)

PIMPINAN UMUM/ PENANGGUNG JAWAB
Dekan Fakultas Matematika dan Ilmu Pengetahuan Alam
Universitas Islam Indonesia

WAKIL PIMPINAN UMUM/ WAKIL PENANGGUNG JAWAB
Ketua Jurusan Farmasi FMIPA UII

MITRA BESTARI

1. Prof. Dr. Wiryatun Lestariana, Apt
2. Prof. Dr. Zullies Ikawati, Apt
3. Prof. Dr. Sudibyo Martono, Apt
4. Dr. Tedjo Yuwono, Apt
5. Prof. Dr. Dachriyanus, Apt
6. Prof. dr. Iwan Dwiprahasto, MMedSc, PhD
7. Prof. Dr. Lukman Hakim M.Sc., Apt
8. Prof. Dr. Achmad Fudholi, DEA, Apt
9. Prof. Dr. Ibnu Gholib Gandjar, DEA., Apt

DEWAN EDITOR

Ketua : Saepudin, M.Si., Apt
Sekretaris : Rochmy Istikharah, M.Sc., Apt.
Anggota : Vitarani Dwi Ananda Ningrum, M.Si., Apt
Okti R. Mafruhah, MSc., Apt
Dimas Adhi Pradana, MSc., Apt.
Fithria DA. Suryanegara, MSc., Apt.
Ari Wibowo, S.Farm., Apt
Arba Pramudita Ramadani, MSc., Apt.
Oktavia Indrati, S.Farm., Apt.

Penerbit

Jurusan Farmasi Fakultas Matematika dan Ilmu Pengetahuan Alam
Universitas Islam Indonesia

Alamat Penerbit

Jurusan Farmasi FMIPA UII
Jl. Kaliurang Km. 14,4 Yogyakarta 55584
Telp. (0274) 896439 ext. 3047
Email: jif@uii.ac.id

PROTECTIVE EFFECTS OF ETHYL ACETATE FRACTION OF *Phaleria macrocarpa* (Scheff) Boerl. ON OXIDATIVE STRESS ASSOCIATED WITH ALLOXAN-INDUCED DIABETIC RATS

Asih Triastuti ^{1*}, Jong Won Choi ²

¹ Department of Pharmacy, Islamic University of Indonesia, Yogyakarta 55584, Indonesia

² College of Pharmacy, Kyungsung University, Busan 608-736, Korea

*e-mail: atea_bening@yahoo.com

ABSTRACT

Oxidative stress resulting from the increased production of reactive oxygen species plays a key role in the development of diabetes mellitus and its complications. *Phaleria macrocarpa*, a traditional plant from Indonesia, has been used empirically to control cancer, arthritis, and diabetes in society. The anti-diabetic effect and antioxidant activity of ethyl acetate (EtOAc) of *Phaleria macrocarpa* (PM) was investigated in alloxan-induced diabetic rats. After two weeks administration of PM, liver antioxidant enzyme and hyperglycemic state were evaluated. The result showed that EtOAc fraction treatments reduced blood glucose levels in diabetic rats by oral administration ($P<0.05$). Serum glutamic-oxaloacetic transaminase (sGOT) and serum glutamic-pyruvate-transaminase (sGPT) also diminished by PM supplementation. The superoxide dismutase (SOD), catalase (CAT), and glutathione-peroxidase (GPx) activities, and glutathione (GSH) level in the alloxan-induced diabetic rats were decreased significantly ($P<0.05$) compared with those in the normal rats but restored by EtOAc fraction treatments. The result of the present study concluded that the diabetic effect of EtOAc fraction of PM against alloxan-induced diabetic rats is mediated either by preventing the decline of hepatic antioxidant status or due to its indirect radical scavenging capacity.

Keywords: Ethyl acetate, Diabetic, Oxidative Stress, *Phaleria macrocarpa*